

STD MORBIDITY

Chlamydia, Gonorrhea and Syphilis Morbidity – January – December 2011

Gender	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Female	20,064	72.2%	3,690	56.2%	13	7.5%
Male	7,682	27.6%	2,867	43.6%	160	92.5%
Unknown	55	<1%	12	<1%	S	S
Total	27,801	100%	6,569	100%	173	100%
Race	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
White	10,757	38.7%	1,726	26.3%	85	49.1%
Black	9,981	35.9%	3,724	56.7%	81	46.8%
Amer. Indian	24	<1%	7	<1%	S	S
Asian	73	<1%	15	<1%	S	S
Multiple Races	9	<1%	S	S	S	S
All Others	1,250	4.5%	214	3.3%	7	4.0%
Unknown	5,707	20.5%	882	13.4%	S	S
Total	27,801	100%	6,569	100%	173	100%
Ethnicity	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Hispanic	999	3.6%	162	2.5%	9	5.2%
Non-Hispanic	15,451	55.6%	4,105	62.5%	157	90.8%
Unknown	11,351	40.8%	2,302	35.0%	7	4.0%
Total	27,801	100%	6,569	100%	173	100%
Age	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
0-9	9	<1%	S	<1%	S	S
10-14	287	1%	49	<1%	S	S
15-19	9,498	34.2%	1,776	27.0%	17	9.8%
20-24	10,813	38.9%	2,488	37.9%	43	24.9%
25-29	3,999	14.4%	1,053	16.0%	38	22.0%
30-34	1,766	6.4%	549	8.4%	21	12.1%
35-39	711	2.6%	291	4.4%	18	10.4%
40-44	376	1.4%	174	2.6%	16	9.2%
45-54	250	<1%	151	2.3%	17	9.8%
55-64	58	<1%	30	<1%	S	S
65+	34	<1%	7	<1%	S	S
Unknown	S	S	S	S	S	S
Total	27,801	100%	6,569	100%	173	100%

Source: ISDH STD Database as of July 6, 2012

NOTE: Not all percents may add to 100 due to rounding. Data is suppressed (S) when counts are <5 or if determined identifiable, by having a corresponding column found to be <5.